

## Flags in zero dimensional complete intersection algebras and indices of real vector fields

L. Giraldo · X. Gómez-Mont · P. Mardešić

Received: 6 December 2006 / Accepted: 4 October 2007  
© Springer-Verlag 2007

- 1 **Abstract** We introduce bilinear forms in a flag in a complete intersection local  $\mathbb{R}$ -algebra  
2 of dimension 0, related to the Eisenbud–Levine, Khimshiashvili bilinear form. We give a  
3 variational interpretation of these forms in terms of Jantzen’s filtration and bilinear forms.  
4 We use the signatures of these forms to compute in the real case the constant relating the GSV-  
5 index with the signature function of vector fields tangent to an even dimensional hypersurface  
6 singularity, one being topologically defined and the other computable with finite dimensional  
7 commutative algebra methods.
- 8 **Keywords** Singularities of functions · Local algebra · Bilinear form · Index of vector field
- 9 **Mathematics Subject Classification (2000)** 58K45 · 58K05 · 13H10

---

Partially supported by Plan Nacional I+D grant no. MTM2004-07203-C02-02, Spain and CONACYT 40329, México.

---

L. Giraldo  
Dep. Matemáticas, Universidad de Cádiz, Puerto Real, Spain

*Present Address:*  
L. Giraldo (✉)  
Dep. Geometría y Topología, F. Matemáticas, Universidad Complutense de Madrid,  
28040 Madrid, Spain  
e-mail: luis.giraldo@uca.es

X. Gómez-Mont  
CIMAT, A.P. 402, 36000 Guanajuato, Mexico  
e-mail: gmont@cimat.mx

P. Mardešić  
Institut de Mathématique de Bourgogne, U.M.R. 5584 du C.N.R.S., Université de Bourgogne,  
B.P. 47 870, 21078 Dijon Cedex, France  
e-mail: mardesic@u-bourgogne.fr

## 10 0 Introduction

11 Let  $f_1, \dots, f_n : \mathbb{R}^n \rightarrow \mathbb{R}$  be germs of real analytic functions that form a regular sequence  
12 as holomorphic functions and let

$$13 \quad \mathbf{A} := \frac{\mathcal{A}_{\mathbb{R}^n,0}}{(f_1, \dots, f_n)} \quad (1)$$

14 be the quotient finite dimensional algebra, where  $\mathcal{A}_{\mathbb{R}^n,0}$  is the algebra of germs of real analytic  
15 functions on  $\mathbb{R}^n$  with coordinates  $x_1, \dots, x_n$ . The class of the Jacobian

$$16 \quad J = \det \left( \frac{\partial f_i}{\partial x_j} \right)_{i,j=1,\dots,n}, \quad J_{\mathbf{A}} := [J]_{\mathbf{A}} \in \mathbf{A} \quad (2)$$

17 generates the socle (the unique minimal non-zero ideal) of the algebra  $\mathbf{A}$ . A symmetric  
18 bilinear form

$$19 \quad \langle \cdot, \cdot \rangle_{L_{\mathbf{A}}} : \mathbf{A} \times \mathbf{A} \xrightarrow{L_{\mathbf{A}}} \mathbb{R} \quad (3)$$

20 is defined by composing multiplication in  $\mathbf{A}$  with any linear map  $L_{\mathbf{A}} : \mathbf{A} \rightarrow \mathbb{R}$  sending  $J_{\mathbf{A}}$   
21 to a positive number. The theory of Eisenbud–Levine and Khimshiashvili asserts that this  
22 bilinear form is nondegenerate and that its signature  $\sigma_{\mathbf{A}}$  is independent of the choice of  $L_{\mathbf{A}}$   
23 (see [3, 12]).

24 Let  $f \in \mathbf{A}$  be an element in the maximal ideal. We define a flag of ideals in  $\mathbf{A}$ :

$$25 \quad K_m := \text{Ann}_{\mathbf{A}}(f) \cap (f^{m-1}), \quad m \geq 1, \quad 0 \subset K_{\ell+1} \subset \dots \subset K_1 \subset K_0 := \mathbf{A} \quad (4)$$

26 and a family of bilinear forms

$$27 \quad \langle \cdot, \cdot \rangle_{L_{\mathbf{A}},f,m} : K_m \times K_m \rightarrow \mathbb{R}, \quad \langle a, a' \rangle_{L_{\mathbf{A}},f,m} = \left\langle \frac{a}{f^{m-1}}, a' \right\rangle_{L_{\mathbf{A}}}, \quad (5)$$

28 defined for  $m = 0, \dots, \ell + 1$ . The division by  $f^{m-1}$  is defined up to elements in  $\text{Ann}_{\mathbf{A}}(f^{m-1})$ ,  
29 but as  $a' \in (f^{m-1})$ , the last expression in (5) is well defined. We call the form  $\langle \cdot, \cdot \rangle_{L_{\mathbf{A}},f,m}$ , the  
30 order  $m$  bilinear form on the algebra  $\mathbf{A}$ , with respect to  $f$ . In Sect. 1 we prove:

31 **Theorem 0.1** For  $m = 0, \dots, \ell + 1$  the order  $m$  bilinear form  $\langle \cdot, \cdot \rangle_{L_{\mathbf{A}},f,m}$  on  $K_m$  induces a  
32 non-degenerate bilinear form

$$33 \quad \langle \cdot, \cdot \rangle_{L_{\mathbf{A}},f,m} : \frac{K_m}{K_{m+1}} \times \frac{K_m}{K_{m+1}} \rightarrow \mathbb{R}, \quad (6)$$

34 whose signature  $\sigma_{\mathbf{A},f,m}$  is independent of the linear map  $L_{\mathbf{A}}$  chosen.

35 In Sect. 2 we give a variational interpretation of Theorem 0.1. Consider germs of analytic  
36 functions  $f, f_1, f_2, \dots, f_n$  in  $\mathbb{R}^n$  such that  $f, f_2, \dots, f_n$  and  $f_1, \dots, f_n$  are regular sequences  
37 as holomorphic functions. We consider the 1-parameter family of ideals  $(f - t, f_2, \dots, f_n)$ .  
38 Choose a small neighborhood  $U_{\mathbb{C}}$  of  $0 \in \mathbb{C}^n$  and a small  $\varepsilon > 0$  such that:

39 (1) The sheaf of algebras on  $U_{\mathbb{C}}$  defined by

$$40 \quad \mathcal{B}_{\mathbb{C}} := \frac{\mathcal{O}_{U_{\mathbb{C}}}}{(f_2, \dots, f_n)}$$

41 is the structure sheaf of a 1-dimensional complete intersection  $\mathbf{Z}_{\mathbb{C}} \subset U_{\mathbb{C}}$  such that the map

$$42 \quad f : \mathbf{Z}_{\mathbb{C}} \rightarrow \Delta_{\varepsilon} \quad (7)$$

43 to the disk  $\Delta_\varepsilon$  of radius  $\varepsilon$  in  $\mathbb{C}$  is a finite analytic map, the sheaf  $f_*\mathcal{B}_\mathbb{C}$  is a free  $\mathcal{O}_{\Delta_\varepsilon}$ -sheaf of  
 44 rank  $\nu$  and  $f^{-1}(0) = 0$ .

45 (2)  $f_1|_{\mathbb{Z}_\mathbb{C}-\{0\}}$  is non-vanishing.

46 These conditions can be fulfilled due to the regular sequence hypothesis [4, 10]. Denoting  
 47 by  $f_*\mathcal{B}^+$  the sheaf on  $(-\varepsilon, \varepsilon) \subset \mathbb{R}$  whose sections are the fixed points of the conjugation  
 48 map  $\bar{\cdot} : f_*\mathcal{B}_\mathbb{C} \rightarrow f_*\mathcal{B}_\mathbb{C}$ , we have that  $f_*\mathcal{B}^+$  is a free  $\mathcal{A}_{(-\varepsilon, \varepsilon)}$ -sheaf of rank  $\nu$ . Its stalk over  
 49  $0$  is

$$50 \quad \mathcal{B} := (f_*\mathcal{B})_0^+ = \frac{\mathcal{A}_{\mathbb{R}^n, 0}}{(f_2, \dots, f_n)}.$$

51 Hence  $\mathcal{B}$  is a free  $\mathcal{A}_{\mathbb{R}, 0}$ -module of rank  $\nu$ . Introduce the 1-parameter family of  $\mathbb{R}$ -algebras  
 52 obtained by evaluation

$$53 \quad \mathbf{B}_{t_0} = f_*\mathcal{B}^+ \otimes_{\mathbb{R}} \frac{\mathbb{R}[t](t-t_0)}{(t-t_0)} = \left[ \bigoplus_{p \in \mathbb{Z}_\mathbb{C} \cap f^{-1}(t_0)} \frac{\mathcal{O}_{\mathbb{C}^n, p}}{(f-t_0, f_2, \dots, f_n)} \right]^+. \quad (8)$$

54  $\mathbf{B}_0$  is a local algebra,  $\mathbf{B}_{t_0}$  is a multilocal algebra and they form a vector bundle of rank  $\nu$  over  
 55  $(-\varepsilon, \varepsilon)$ , whose sheaf of real analytic sections is  $f_*\mathcal{B}^+$ .

56 We define in the sheaf of sections  $f_*\mathcal{B}^+$ , a bilinear map

$$57 \quad \langle \cdot, \cdot \rangle : f_*\mathcal{B}^+ \times f_*\mathcal{B}^+ \rightarrow f_*\mathcal{B}^+ \xrightarrow{\mathcal{L}} \mathcal{A}_{(-\varepsilon, \varepsilon)}, \quad \langle a, b \rangle = \mathcal{L}(a \cdot b),$$

58 obtained by first applying the multiplication in the sheaf of algebras  $f_*\mathcal{B}^+$  and then applying  
 59 a chosen  $\mathcal{A}_{(-\varepsilon, \varepsilon)}$ -module map  $\mathcal{L} : f_*\mathcal{B}^+ \rightarrow \mathcal{A}_{(-\varepsilon, \varepsilon)}$  having the property that evaluating it  
 60 at  $0$  gives a linear map  $L_{\mathbf{B}_0} : \mathbf{B}_0 \rightarrow \mathbb{R}$ , verifying  $L_{\mathbf{B}_0}([J]_{\mathbf{B}_0}) > 0$ . The evaluation of  $\langle \cdot, \cdot \rangle$   
 61 at a fiber  $\mathbf{B}_t$  is a bilinear form defined on  $\mathbf{B}_t$  and denoted by  $\langle \cdot, \cdot \rangle_t$ .

62 This family of non-degenerate bilinear forms is the usual tool in the Eisenbud–Levine  
 63 and Khimshiashvili theory [3, 12] to calculate the degree of the smooth map given by  
 64  $(f, f_2, \dots, f_n) : (\mathbb{R}^n, 0) \rightarrow (\mathbb{R}^n, 0)$ .

65 Define a sheaf map by multiplication with  $f_1$

$$66 \quad M_{f_1} : f_*\mathcal{B}^+ \rightarrow f_*\mathcal{B}^+ \quad M_{f_1}(b) = f_1 b$$

67 and a family of bilinear maps, that we call relative:

$$68 \quad \langle \cdot, \cdot \rangle^{rel} : f_*\mathcal{B}^+ \times f_*\mathcal{B}^+ \rightarrow \mathcal{A}_{(-\varepsilon, \varepsilon)}, \quad \langle a, b \rangle^{rel} = \langle M_{f_1}(a), b \rangle \quad (9)$$

$$69 \quad \langle \cdot, \cdot \rangle_t^{rel} : \mathbf{B}_t \times \mathbf{B}_t \rightarrow \mathbb{R}, \quad \langle [a]_t, [b]_t \rangle_t^{rel} = \langle M_{[f_1]_t}([a]_t), [b]_t \rangle_t. \quad (10)$$

70 The bilinear forms  $\langle \cdot, \cdot \rangle_t^{rel}$  are non-degenerate, for  $t \neq 0$ , having signature  $\tau_\pm$ , for  $\pm t > 0$ .  
 71 The form  $\langle \cdot, \cdot \rangle_t^{rel}$  degenerates for  $t = 0$  on  $\text{Ann}_{\mathbf{B}_0}([f_1]_{\mathbf{B}_0})$  [5]. Expanding in Taylor series at  
 72  $0$  the family of relative bilinear forms we arrive at the setting in [11, 14], where it is shown  
 73 how to obtain a flag of ideals

$$74 \quad \dots \subset \tilde{K}_r \subset \dots \subset \tilde{K}_1 \subset \tilde{K}_0 = \mathbf{B}_0 \quad (11)$$

75 and bilinear forms in them and show how to reconstruct from the signatures  $\tau_m$  of these  
 76 bilinear forms the signatures  $\tau_\pm$  (see Proposition 2.1). In our algebraic setting, the flag and  
 77 the bilinear forms have the algebraic description.

78 **Theorem 0.2** For the family of bilinear forms  $\langle \cdot, \cdot \rangle_t^{rel}$ , in the family of algebras  $\mathbf{B}_t$  (10) we  
 79 have

- 80 (1) The set of  $b \in \mathcal{B}$  such that the function  $t \rightarrow \langle [b]_t, [b']_t \rangle_t^{rel}$  vanishes at 0 up to order  $m$ ,  
81 for every  $b' \in \mathcal{B}$  is the quotient ideal

$$(f^m : f_1) := \{b \in \mathcal{B} / f_1 b \in (f^m)\} \subset \mathcal{B}$$

82 and

$$\tilde{K}_m = \frac{(f^m : f_1)}{(f) \cap (f^m : f_1)} \subset \frac{\mathcal{B}}{(f)} = \mathbf{B}_0. \quad (12)$$

- 85 (2)  $(f) \cap (f^m : f_1) = M_f((f^{m-1} : f_1))$ .

- 86 (3) The bilinear form  $(b, b') \rightarrow L_{\mathbf{B}_0} \left( \left[ \begin{smallmatrix} f_1 b \\ f^m b' \end{smallmatrix} \right]_{\mathbf{B}_0} \right)$

$$(f^m : f_1) \oplus (f^m : f_1) \xrightarrow{\frac{f_1}{f^m}} (f^m : f_1) \xrightarrow{\tilde{\pi}_0} \frac{\mathcal{B}}{(f)} = \mathbf{B}_0 \xrightarrow{L_{\mathbf{B}_0}} \mathbb{R}, \quad (13)$$

88 where  $\tilde{\pi}_0$  is the projection from  $\mathcal{B}$  to  $\mathcal{B}/(f) = \mathbf{B}_0$ , vanishes on  $(f) \cap (f^m : f_1)$  and  
89 induces Jantzen's bilinear form

$$\langle \cdot, \cdot \rangle^m : \tilde{K}_m \otimes \tilde{K}_m \rightarrow \mathbb{R} \quad \langle \cdot, \cdot \rangle^m = \left\langle \frac{f_1}{f^m}, \cdot \right\rangle_0, \quad (14)$$

91 giving the formula

$$\tau_+ = \sum_{m \geq 0} \tau_m, \quad \tau_- = \sum_{m \geq 0} (-1)^m \tau_m$$

93 In Sect. 3, we show

94 **Theorem 0.3** There is an isomorphism  $\varphi : \tilde{K}_1 \rightarrow K_1$ , induced by multiplication with the  
95 function  $\frac{f_1}{f}$ , which is sending the flag  $\{\tilde{K}_m\}_{m \geq 1}$  in  $\mathbf{B}_0$  in (12) to the flag  $\{K_m\}_{m \geq 1}$  in  $\mathbf{A}$  in (4)  
96 and Jantzen's bilinear forms (14) to the bilinear forms (6). Hence, for  $m \geq 1$ , we have equal  
97 signatures  $\tau_m = \sigma_{\mathbf{A}, f, m}$  and

$$\tau_+ = \tau_0 + \sum_{m=1}^{\ell+1} \sigma_{\mathbf{A}, f, m}, \quad \tau_- = \tau_0 + \sum_{m=1}^{\ell+1} (-1)^m \sigma_{\mathbf{A}, f, m}.$$

99 In Sect. 4 we apply these considerations for calculating indices of vector fields. If  $X =$   
100  $\sum_{i=1}^n X^i \frac{\partial}{\partial x_i}$  is a real analytic vector field with an algebraically isolated zero at 0 in  $\mathbb{R}^n$ , then  
101 the (Poincaré-Hopf) index of  $X$  at 0 is the signature of the bilinear form (3) constructed for  
102 the finite dimensional algebra

$$\mathbf{B} := \frac{\mathcal{A}_{\mathbb{R}^n, 0}}{(X^1, \dots, X^n)}, \quad \langle \cdot, \cdot \rangle_{L_{\mathbf{B}}} : \mathbf{B} \times \mathbf{B} \rightarrow \mathbf{B} \xrightarrow{L_{\mathbf{B}}} \mathbb{R}$$

104 where  $L_{\mathbf{B}} : \mathbf{B} \rightarrow \mathbb{R}$  is a linear map with  $L_{\mathbf{B}}(J_X) > 0$  (see [3, 12]). Now assume further that  
105  $f : (\mathbb{R}^n, 0) \rightarrow (\mathbb{R}, 0)$  is a real analytic function, that  $X$  is tangent to the fiber  $V_0 := f^{-1}(0)$ ,  
106 giving the relation  $df(X) = hf$  with  $h$  a real analytic function called the cofactor. If 0 is a  
107 smooth point of  $V_0$  then the signature  $\sigma_{\mathbf{B}, h, 0}$  of the order 0 bilinear form

$$\langle \cdot, \cdot \rangle_{L, h, 0} : \frac{\mathbf{B}}{\text{Ann}_{\mathbf{B}}(h)} \times \frac{\mathbf{B}_{\mathbb{R}}}{\text{Ann}_{\mathbf{B}}(h)} \rightarrow \frac{\mathbf{B}}{\text{Ann}_{\mathbf{B}}(h)} \xrightarrow{L} \mathbb{R}$$

$$L : \frac{\mathbf{B}_{\mathbb{R}}}{\text{Ann}_{\mathbf{B}}(h)} \rightarrow \mathbb{R}, \quad L \left( \frac{J_{\mathbf{B}}}{h} \right) > 0$$

110 is the Poincaré-Hopf index at 0 of the vector field  $X|_{V_0}$ , as can easily be deduced using the  
 111 implicit function theorem. If 0 is an isolated critical point of  $V_0$  and the dimension  $n$  of the  
 112 ambient space is even, in [7] it is proved that

$$113 \quad \text{Ind}_{V_{+,0}}(X) = \text{Ind}_{V_{-,0}}(X) = \sigma_{\mathbf{B},h,0} - \sigma_{\mathbf{A},h,0}. \quad (15)$$

114 If  $n$  is odd, it is proved in [6] that

$$115 \quad \text{Ind}_{V_{\pm,0}}(X) = \sigma_{\mathbf{B},h,0} + K_{\pm}. \quad (16)$$

116 In the case of odd dimensional ambient space and for  $f$  a germ of a real analytic function  
 117 with an algebraically isolated singularity at 0, we calculate the constants  $K_{\pm}$  by studying the  
 118 family of contact vector fields

$$119 \quad X_t = (f - t) \frac{\partial}{\partial x_1} + \sum_{i=1}^N \left[ \frac{\partial f}{\partial x_{2i+1}} \frac{\partial}{\partial x_{2i}} - \frac{\partial f}{\partial x_{2i}} \frac{\partial}{\partial x_{2i+1}} \right],$$

120 where  $f_j := \frac{\partial f}{\partial x_j}$ . For  $t \neq 0$ , the signatures of the relative bilinear forms correspond to  
 121 the sum of the Poincaré-Hopf indices of the restriction of  $X_t$  to  $V_t$ . Our transport from the  
 122 algebra  $\mathbf{B}_0$  to the Jacobian algebra  $\mathbf{A}$  is a local analogue of the Poincaré-Hopf Theorem  
 123 relating information of the singular point of  $X$  to invariants of the singularity of  $f$ .

124 Using these explicit computations for contact vector fields, we conclude the search for an  
 125 algebraic formula for the real GSV-index using local algebra by determining the values of  
 126 the constants  $K_{\pm}$ :

127 **Theorem 0.4** *Let  $V$  be an algebraically isolated hypersurface singularity in  $\mathbb{R}^{2N+1}$ , then*  
 128 *the constants  $K_{\pm}$  in (16) relating the GSV-index and the signature  $\sigma_{\mathbf{B},h,0}$  are:*

$$129 \quad K_+ = \sum_{m \geq 1} \sigma_{\mathbf{A},f,m}, \quad K_- = \sum_{m \geq 1} (-1)^m \sigma_{\mathbf{A},f,m}.$$

### 130 1 Higher order signatures in $\mathbf{A}$

131 We use the definitions and notations of Sect. 0. The algebra  $\mathbf{A}$  has an intrinsic  $\mathbf{A}$ -valued  
 132 bilinear map, which is the multiplication in  $\mathbf{A}$ :

$$133 \quad (\cdot, \cdot)_{\mathbf{A}} : \mathbf{A} \times \mathbf{A} \longrightarrow \mathbf{A} \quad (a, b)_{\mathbf{A}} := ab. \quad (17)$$

134 In terms of this pairing and the non-singular pairing  $\langle \cdot, \cdot \rangle_{L_{\mathbf{A}}}$  in (3), the orthogonal of an ideal  
 135  $I \subset \mathbf{A}$  is the annihilator ideal in the algebra  $\mathbf{A}$ :  $I^{\perp} = \text{Ann}_{\mathbf{A}}(I)$ . The process of taking the  
 136 orthogonal induces an involution in the set of ideals of  $\mathbf{A}$ , which is reversing the natural  
 137 inclusions of sets in  $\mathbf{A}$ . In particular, the orthogonal to the maximal ideal is the socle. It is  
 138 1-dimensional and the class of the Jacobian  $J_{\mathbf{A}}$  is a generator (see [3,4,12]).

139 Choose now an element  $f \in \mathbf{A}$  in the maximal ideal. Consider the linear map induced in  
 140  $\mathbf{A}$  by multiplication with  $f$ :

$$141 \quad M_f : \mathbf{A} \longrightarrow \mathbf{A} \quad M_f(a) = fa.$$

142 For  $j \geq 1$ , the maps  $M_f^j$  are selfadjoint maps for the bilinear map (17):

$$143 \quad (M_f^j a, b)_{\mathbf{A}} = f^j ab = af^j b = (a, M_f^j b)_{\mathbf{A}},$$

144 and hence they are also selfadjoint maps for the bilinear form  $\langle \cdot, \cdot \rangle_{L_{\mathbf{A}}}$ . We have that

$$145 \quad \text{Ann}_{\mathbf{A}}(f^j) = \text{Ker}(M_f^j) \quad \text{and} \quad (f^j) = \text{Im}(M_f^j)$$

146 and each of these spaces is the orthogonal of the other in  $\mathbf{A}$ , since  $M_f^j$  is selfadjoint.

147 Consider the flag of ideals in  $\mathbf{A}$

$$148 \quad 0 \subset (f^\ell) \subset (f^{\ell-1}) \subset \dots \subset (f^2) \subset (f) \subset \mathbf{A}, \quad (18)$$

149 where  $\ell$  is minimal with  $f^{\ell+1} = 0$  and the orthogonal flag of ideals

$$150 \quad 0 \subset \text{Ann}_{\mathbf{A}}(f) \subset \text{Ann}_{\mathbf{A}}(f^2) \subset \dots \subset \text{Ann}_{\mathbf{A}}(f^{\ell-1}) \subset \text{Ann}_{\mathbf{A}}(f^\ell) \subset \mathbf{A}. \quad (19)$$

151 The linear map  $M_f : \mathbf{A} \rightarrow \mathbf{A}$  is a nilpotent map  $M_f^{\ell+1} = 0$ .

152 **Lemma 1.1** For  $j = 1, \dots, \ell + 1$ , there are linear subspaces  $P_j$  of  $\mathbf{A}$ , called primitive  
153 subspaces, such that

$$154 \quad \mathbf{A} = \bigoplus_{j=1}^{\ell+1} \left[ \bigoplus_{k=0}^{j-1} M_f^k P_j \right], \quad (20)$$

155 with  $M_f^{j-1} : P_j \rightarrow \mathbf{A}$  injective and  $M_f^j(P_j) = 0$ . The mapping  $M_f : \mathbf{A} \rightarrow \mathbf{A}$  is in  
156 Jordan canonical form in any basis obtained by choosing bases of each of the spaces  $P_j$  and  
157 extending them to a basis of  $\mathbf{A}$  by the action of  $M_f$  as in (20).

158 *Proof* We recall how to choose a basis of  $\mathbf{A}$  as a vector space over  $\mathbb{R}$  that expresses  $M_f$  in  
159 Jordan canonical form. Inductively, let us begin by choosing linearly independent vectors  
160  $v_1, \dots, v_{n_{\ell+1}}$  generating a vector space  $P_{\ell+1}$  complementary to  $\text{Ann}_{\mathbf{A}}(f^\ell)$  in  $\mathbf{A}$  and choose  
161 as first vectors of a basis of  $\mathbf{A}$  the vectors

$$162 \quad \{v_j, f v_j, \dots, f^\ell v_j\}_{j=1, \dots, n_{\ell+1}}.$$

163 With  $P_{\ell+1}$  we construct the Jordan blocks of maximal size  $\ell$  of  $M_f$ . Then, we choose linearly  
164 independent vectors  $v_{n_{\ell+1}+1}, \dots, v_{n_{\ell+1}+n_\ell}$  generating a vector space  $P_\ell$  with the property  
165 that

$$166 \quad \text{Ann}_{\mathbf{A}}(f^{\ell-1}) \oplus M_f(P_{\ell+1}) \oplus P_\ell = \text{Ann}_{\mathbf{A}}(f^\ell).$$

167 We choose the next part of the basis by choosing the vectors

$$168 \quad \{v_j, f v_j, \dots, f^{\ell-1} v_j\}_{j=n_{\ell+1}+1, \dots, n_{\ell+1}+n_\ell}$$

169 to construct the Jordan blocks of size  $\ell - 1$ , and so on. The space of 1-st primitive vectors  
170  $P_1$  is formed of vectors in  $\mathbf{A}$  with the property that

$$171 \quad M_f^\ell(P_{\ell+1}) \oplus M_f^{\ell-1}(P_\ell) \oplus \dots \oplus M_f^2(P_3) \oplus M_f(P_2) \oplus P_1 = \text{Ann}_{\mathbf{A}}(f).$$

172 □

173 We call the vectors in  $P_j$   $j$ th-primitive vectors, and we denote by  $n_j$  the dimension of  $P_j$ .  
174 Hence  $n_j$  is also the number of Jordan blocks of size  $j$  in  $M_f$ . It is convenient to present the

175 direct sum decomposition (20) by the matrix:

$$176 \quad \mathbf{A} = \begin{pmatrix} P_1 & P_2 & P_3 & P_4 & \cdots & P_\ell & P_{\ell+1} \\ 0 & M_f P_2 & M_f P_3 & M_f P_4 & \cdots & M_f P_\ell & M_f P_{\ell+1} \\ 0 & 0 & M_f^2 P_3 & M_f^2 P_4 & \cdots & M_f^2 P_\ell & M_f^2 P_{\ell+1} \\ & & & & \cdots & & \\ 0 & 0 & 0 & 0 & \cdots & M_f^{\ell-1} P_\ell & M_f^{\ell-1} P_{\ell+1} \\ 0 & 0 & 0 & 0 & \cdots & 0 & M_f^\ell P_{\ell+1} \end{pmatrix}, \quad (21)$$

177 meaning that an element of  $\mathbf{A}$  has components in the form of an upper triangular matrix where  
 178 the  $(i, j)$ th-entry of the matrix is an arbitrary element in  $M_f^{i-1}(P_j)$ , with  $i, j = 1, \dots, \ell + 1$ .  
 179 Each column is formed by equidimensional subspaces, until we reach the zero subspace, and  
 180 the map  $M_f$  acts as a map preserving columns and descending one row. Hence, restricting  
 181 to a column in (21), the map  $M_f$  is an isomorphism until it reaches the diagonal, where  $M_f$   
 182 is the zero map.

183 Using this representation for  $\mathbf{A}$  and recalling the flag of ideals (4), we have

- 184 **Lemma 1.2** (1) *The ideal  $(f^m)$  is formed by the last  $\ell + 1 - m$  rows of the matrix (21).*  
 185 (2) *Its orthogonal  $\text{Ann}_{\mathbf{A}}(f^m)$  is formed by the elements in a band of width  $m$  above the*  
 186 *diagonal in (21), including the diagonal.*  
 187 (3) *The ideal  $K_m$  in (4) is formed by the lower  $\ell + 2 - m$  diagonal terms.*  
 188 (4) *The ideal  $K_m^\perp$ , orthogonal to  $K_m$  is*

$$189 \quad K_m^\perp = (f) + \text{Ann}_{\mathbf{A}}(f^{m-1}). \quad (22)$$

190 *Example 1.1* For  $\ell = 3$  and  $m = 3$ , we have

$$191 \quad (f^2) = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & M_f^2 P_3 & M_f^2 P_4 \\ 0 & 0 & 0 & M_f^3 P_4 \end{pmatrix} \quad \text{Ann}_{\mathbf{A}}(f^2) = \begin{pmatrix} P_1 & P_2 & 0 & 0 \\ 0 & M_f P_2 & M_f P_3 & 0 \\ 0 & 0 & M_f^2 P_3 & M_f^2 P_4 \\ 0 & 0 & 0 & M_f^3 P_4 \end{pmatrix}$$

$$192 \quad K_3 = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & M_f^2 P_3 & 0 \\ 0 & 0 & 0 & M_f^3 P_4 \end{pmatrix}; \quad (f) + \text{Ann}_{\mathbf{A}}(f^2) = \begin{pmatrix} P_1 & P_2 & 0 & 0 \\ 0 & M_f P_2 & M_f P_3 & M_f P_4 \\ 0 & 0 & M_f^2 P_3 & M_f^2 P_4 \\ 0 & 0 & 0 & M_f^3 P_4 \end{pmatrix} = K_3^\perp.$$

193 *Proof of Lemma 1.2* Since  $M_f$  corresponds to going down 1 row in (21), parts 1, 2 and 3,  
 194 are clear. To prove part 4, note first that

$$195 \quad (f) + \text{Ann}_{\mathbf{A}}(f^{m-1}) \subset K_m^\perp.$$

196 The ideal  $(f)$  is given by all the terms in (21), except for the first row. Since  $\text{Ann}_{\mathbf{A}}(f^{m-1})$  is  
 197 the band matrix above the diagonal of width  $m - 1$ , we obtain that the only contribution of  
 198  $(f) + \text{Ann}_{\mathbf{A}}(f^{m-1})$  to  $(f)$  is given by the first  $m - 1$  terms in the first row. On the other hand  
 199  $K_m = \text{Ann}_{\mathbf{A}}(f) \cap (f^{m-1})$  consist of the last  $\ell + 2 - m$  terms in the diagonal. We observe  
 200 on using (21) that the ideals  $(f) + \text{Ann}_{\mathbf{A}}(f^{m-1})$  and  $K_m$  have complementary dimensions  
 201 in  $\mathbf{A}$ . Now (22) must hold, as the bilinear form  $\langle \cdot, \cdot \rangle_{L_{\mathbf{A}}}$  is non-degenerate.  $\square$

202 **Proposition 1.1** For the bilinear forms in (5), we have

- 203 (1)  $\langle \cdot, \cdot \rangle_{L_A, f, 0} = \langle f \cdot, \cdot \rangle_{L_A}$  has  $K_1 = \text{Ann}_A(f)$  as degeneracy locus and the induced non-  
 204 degenerate bilinear form in  $\frac{A}{\text{Ann}_A(f)}$  is obtained by choosing  $\frac{J_A}{f}$  as generator of the 1  
 205 dimensional socle of  $\frac{A}{\text{Ann}_A(f)}$  and defining the bilinear form as multiplication followed  
 206 by a real valued map sending  $\frac{J_A}{f}$  to a positive number.  
 207 (2) The bilinear form  $\langle \cdot, \cdot \rangle_{L_A, f, 1} = \langle \cdot, \cdot \rangle_{L_A|_{K_1 \times K_1}}$  has  $K_2$  as degeneracy locus.  
 208 (3) For  $m \geq 2$  the bilinear form  $\langle \cdot, \cdot \rangle_{L_A, f, m}$  in (5) is well defined and has  $K_{m+1}$  as degeneracy  
 209 locus.

210 *Proof* (1) The inner product  $\langle f \cdot, \cdot \rangle_{L_A}$  vanishes on  $\text{Ann}_A(f)$ . If  $\langle fa, a' \rangle_{L_A} = 0$  for all  $a'$ ,  
 211 then  $fa = 0$ , since (3) is a non-degenerate bilinear form on  $A$  [3,4,12]. Hence,  $\langle \cdot, \cdot \rangle_{L_A, f, 0}$  has  
 212  $K_1$  as degeneracy locus. The algebra  $\frac{A}{\text{Ann}_A(f)}$  has a one-dimensional socle generated by the  
 213 class of  $J_A/f$  (see [5] for more details).

214 (2) Note first that  $K_2 = \text{Ann}_A(f) \cap (f) = K_1 \cap K_1^\perp$ , by (4) and (22). Hence, given  $a \in K_2$   
 215 and any  $b \in K_1$ , it follows that  $(a, b)_A = 0$ , so  $K_2$  is contained in the degeneracy locus of  
 216  $\langle \cdot, \cdot \rangle_{L_A, f, 1}$ . On the other hand, let  $a \in K_1 - K_2 = K_1 - K_1^\perp$ . Then  $aK_1$  is a non-zero ideal in  $A$ ,  
 217 and so contains the socle of  $A$ . We obtain an expression  $J_A = ac$ , for some  $c \in K_1$ . Hence,  
 218  $\langle a, c \rangle_{L_A} = L_A(ac) = L_A(J_A) > 0$ , so that  $a$  is not in the degeneracy locus of  $\langle \cdot, \cdot \rangle_{L_A, f, 1}$ .

219 (3) Let  $m \geq 2$ . We first show that the bilinear form  $\langle \cdot, \cdot \rangle_{L_A, f, m}$  is well defined, i.e. is  
 220 independent of the division by  $f^{m-1}$  in  $K_m$ . Let  $a, b$  be in  $K_m = \text{Ann}_A(f) \cap (f^{m-1})$ . Then  
 221 there exists  $a_1 \in A$  such that  $a = a_1 f^{m-1}$  and  $\langle a, b \rangle_{L_A, f, m} = \langle a_1, b \rangle_{L_A}$ . Let also  $a =$   
 222  $a_2 f^{m-1}$ . Then  $\langle a_1, b \rangle_{L_A} = \langle a_2, b \rangle_{L_A}$ , because  $a_1 - a_2 \in \text{Ann}_A(f^{m-1})$  and  $b \in (f^{m-1})$ .

223 If  $a \in K_{m+1}$ , then  $\frac{a}{f^{m-1}} \in (f)$ , and since  $b \in K_m \subset \text{Ann}_A(f)$ , we have  $\frac{a}{f^{m-1}} b = 0$ .  
 224 Hence, the form  $\langle \cdot, \cdot \rangle_{L_A, f, m}$  degenerates on  $K_{m+1}$ .

225 Let  $a \in K_m - K_{m+1}$ . In order to prove that the form  $\langle \cdot, \cdot \rangle_{L_A, f, m}$  is non-degenerate on  $a$ ,  
 226 we have to show that  $\frac{a}{f^{m-1}} \notin K_m^\perp$ . Using the representation (21), and part (3) of Lemma 1.2,  
 227 the  $a_{m,m}$  entry in  $a$  is not zero, and  $a_{m,m} \in M_f^{m-1} P_m$ . Now  $\frac{a}{f^{m-1}}$  is obtained by lifting all  
 228 the elements in the representation by  $m-1$  rows, keeping the columns fixed. We observe  
 229 that  $\frac{a}{f^{m-1}} \notin (f)$ . It now suffices to show that  $\frac{a}{f^{m-1}} \notin \text{Ann}_A(f^{m-1})$ . But by part (4) of  
 230 Lemma 1.2, the space  $\text{Ann}_A(f^{m-1})$  is given by the band matrix of width  $m-1$ , including  
 231 the diagonal. Hence,  $\frac{a}{f^{m-1}}$  is not an element of  $\text{Ann}_A(f^{m-1})$ .  $\square$

232 *Proof of Theorem 0.1* By Proposition 1.1, we have that  $K_{m+1}$  is the locus of the bilinear  
 233 form  $\langle \cdot, \cdot \rangle_{L_A, f, m}$ , so that  $K_m/K_{m+1}$  inherits a non-degenerate bilinear form. The linear forms  
 234  $L_A$ , verifying  $L_A(J_A) > 0$  form an open connected set in the dual space  $\mathbb{R}^{n^*}$ . The signature  
 235 is an integer valued continuous function of  $L_A$ , hence it is constant.  $\square$

236 **Corollary 1** For  $m \geq 1$ , the mapping

$$237 \quad M_f^{m-1} : P_m \longrightarrow K_m/K_{m+1} \quad (23)$$

238 is a well defined isomorphism. The pairing of  $m$ -primitive vectors

$$239 \quad \langle \cdot, \cdot \rangle_{L_A, m}^{prim} : P_m \times P_m \longrightarrow \mathbb{R}, \quad \langle a, b \rangle_{L_A, m}^{prim} := \langle M_f^{m-1} a, b \rangle_{L_A}$$

240 is a non-degenerate symmetric bilinear pairing, induced by the pairing (6) via the isomor-  
 241 phism (23).

242 *Proof* Using the representation (21) for the elements of  $A$  and the description of  $K_m$  given  
 243 in part 3 of Lemma 1.2, we have that  $M_f^{m-1} : P_m \longrightarrow K_m$  is injective, and hence (23) is a

244 well defined isomorphism. The pull back of the non-degenerate bilinear form on  $K_m/K_{m+1}$   
 245 via this last map is

$$\begin{aligned}
 \langle b, b' \rangle_{L_{\mathbf{A}}, m} &= L_{\mathbf{A}}(f^{m-1} \cdot b \cdot b') = L_{\mathbf{A}} \left( \left( \frac{1}{f^{m-1}} \cdot f^{m-1} b \right) \cdot f^{m-1} b' \right) \\
 &= \langle M_f^{m-1}(b), M_f^{m-1}(b') \rangle_{L_{\mathbf{A}}, f, m}.
 \end{aligned}$$

248 □

249 *Example 1.2* Let  $f$  be a germ of a real analytic function in  $\mathbb{R}^n$ , with an algebraically isolated  
 250 critical point. This means that the ideal generated by the partial derivatives of  $f$  in the ring of  
 251 germs of holomorphic functions has finite codimension. Let  $\mathbf{A} = \mathbf{A}(f)$  be given by (1), with  
 252  $f_i := \frac{\partial f}{\partial x_i}$  and let  $\ell$  be as in (18). Let  $\sigma_{f,m} = \sigma_{\mathbf{A}(f), f, m}$ ,  $m = 0, \dots, \ell + 1$ , be the signatures  
 253 given by Theorem 0.1. These are invariants associated to the germ  $f$ . We call  $\sigma_{f,m}$  the order  
 254  $m$  signature of  $f$ .

255 **2 The family of bilinear forms in  $\mathcal{B}$**

256 In this section we construct a family of bilinear forms  $\langle, \rangle^{rel}$ , that we call relative, which is  
 257 constructed from the equations

$$258 \quad f - t = f_2 = \dots = f_n = 0$$

259 which are non-degenerate for  $t \neq 0$ . We do Taylor series expansion of  $\langle, \rangle^{rel}$  and determine  
 260 an algebraic procedure to compute the signatures for  $t \neq 0$  in terms of local linear algebra  
 261 in the ring  $\frac{\mathcal{A}_{\mathbb{R}^n, 0}}{(f_2, \dots, f_n)}$  via the first terms of the above Taylor series expansion.

262 Recall the setting and definitions of Sect. 0. Note in particular that since the map in (7) is a  
 263 finite analytic map, the inverse image  $f^{-1}((-\varepsilon, \varepsilon))$  is a finite union of curves (parameterized  
 264 by  $(-\varepsilon, 0]$  or  $[0, \varepsilon)$ ), which come together at 0. The conjugation map permutes them, and  
 265 the fixed components correspond to  $\mathbf{Z} := \mathbf{Z}_{\mathbb{C}} \cap \mathbb{R}^n$ . Hence  $\mathbf{Z}$  consists either of 0 only or of a  
 266 finite number of these real curves all passing through 0, which is its only singular point. Note  
 267 that the degree of the covering map  $f : \mathbf{Z} - \{0\} \rightarrow (-\varepsilon, \varepsilon) - \{0\}$  may be distinct for  $t > 0$   
 268 and  $t < 0$ . In the sheaf  $f_*\mathcal{B}^+$  we have information about the points  $\{f = t\}_{t \in (-\varepsilon, \varepsilon)} \cap \mathbf{Z}_{\mathbb{C}}$   
 269  $U_{\mathbb{C}}$ , real or complex.

270 **Lemma 2.1** *The signature of the non-degenerate bilinear forms  $\langle, \rangle_t$  on  $\mathbf{B}_t$  is independent*  
 271 *of  $t$  and it is equal to the sum of the signatures of the bilinear forms computed on the local*  
 272 *rings  $\mathbf{B}_{t,p}$  for  $p \in \mathbf{Z} \cap f^{-1}(t)$ , for each  $t \in (-\varepsilon, \varepsilon)$ .*

273 *Proof* This is the usual procedure due to Eisenbud–Levine and Khimshiashvili [3,12]  
 274 to calculate the degree applied to the smooth map given by  $(f, f_2, \dots, f_n) : (\mathbb{R}^n, 0) \rightarrow$   
 275  $(\mathbb{R}^n, 0)$ . In particular, the contribution to the signature coming from points in  $\mathbf{Z}_{\mathbb{C}} - \mathbf{Z}$   
 276 is always 0. □

277 Using a trivialization of  $f_*\mathcal{B}^+$ , we can transfer the relative forms  $\langle, \rangle_t^{rel}$  from  $\mathbf{B}_t$  to  $\mathbf{B}_0$ . So  
 278 we have a family of bilinear forms that we denote by  $\langle, \rangle_t$ . We are interested in the Taylor  
 279 expansion of this family of bilinear forms at  $t = 0$ . We will use the following Proposition,  
 280 containing results from [11,14]:

281 **Proposition 2.1** *Let  $\langle, \rangle_t$ ,  $t \in (-\varepsilon, \varepsilon)$ , be an analytic family of forms on a finite dimensional*  
 282 *vector space  $\mathbf{B}_0$ . Assume that the forms  $\langle, \rangle_t$  are nondegenerate for  $t \neq 0$ . Let  $\tilde{K}_i, i = 0, \dots, r$ ,*

283 be the set of  $[b]_0 \in \mathbf{B}_0$ , such that the functions  $t \mapsto \langle [b]_0, [b']_0 \rangle_t$  vanish at 0 up to order  $i$   
 284 for any  $b' \in \mathcal{B}$ . Then

285 (1) For  $i = 0, \dots, r$ , a bilinear form  $\langle \cdot, \cdot \rangle^i$ , is well defined on  $\tilde{K}_i$  by

286 
$$\langle [b]_0, [b']_0 \rangle^i = \frac{1}{i!} \frac{d^i}{dt^i} \langle b, b' \rangle_t |_{t=0}. \quad (24)$$

287 (2) The bilinear form  $\langle \cdot, \cdot \rangle^i$  degenerates on  $\tilde{K}_{i+1}$ , and induces a nondegenerate bilinear form  
 288 on  $\tilde{K}_i / \tilde{K}_{i+1}$ . Denote its signature by  $\tau_i$ .

289 (3) The signatures  $\tau_+$  and  $\tau_-$  of the forms  $\langle \cdot, \cdot \rangle_t$  on  $\mathbf{B}_0$ , for  $t > 0$  and  $t < 0$ , respectively,  
 290 are given by

291 
$$\tau_+ = \sum_{i=0}^r \tau_i, \quad \tau_- = \sum_{i=0}^r (-1)^i \tau_i. \quad (25)$$

292 *Proof of Theorem 0.2* (1) Let  $b \in \mathcal{B} - \{0\}$ . Then, there is a unique integer  $j$  and  $c \in \mathcal{B}$  with  
 293  $[c]_{\mathbf{B}_0} \neq 0$  such that  $f_1 b = f^j c$ . Since  $[c]_{\mathbf{B}_0} \neq 0$ , and  $[J]_{\mathbf{B}_0}$  is a generator of the socle of  $\mathbf{B}_0$   
 294 we may find  $e_0 \in \mathbf{B}_0$  such that  $[c]_{\mathbf{B}_0} e_0 = [J]_{\mathbf{B}_0}$ . Choose any  $e \in \mathcal{B}$  with the property that  
 295  $[e]_{\mathbf{B}_0} = e_0$ , so that  $\mathcal{L}(ce)(0) = L_{\mathbf{B}_0}([c]_{\mathbf{B}_0} e_0) = L_{\mathbf{B}_0}([J]_{\mathbf{B}_0}) \neq 0$ . For any  $b' \in \mathcal{B}$ , we have  
 296  $f_1 b b' = f^j c b' \in (f^j)$ . Hence,

297 
$$\langle b, b' \rangle^{rel} = \mathcal{L}(f_1 b b') = \mathcal{L}(f^j c b') = t^j \mathcal{L}(c b') \in (t^j)$$

298 and

299 
$$\langle b, e \rangle^{rel} = \mathcal{L}(f_1 b e) = \mathcal{L}(f^j c e) = t^j \mathcal{L}(c e) \in (t^j) - (t^{j+1}).$$

300 Hence, if  $b$  is as in the statement of part (1), we have that  $j \geq m$  and hence  $f_1 b \in (f^m)$ , i.e.  
 301  $b \in (f^m : f_1)$ . This proves the first assertion. The second assertion follows from the first by  
 302 evaluating it at  $t = 0$  and using (8).

303 (2) Let  $b \in (f) \cap (f^m : f_1)$ . Then  $b = cf$  and  $(cf) f_1 = ef^m$ , so that  $cf_1 = ef^{m-1}$ .  
 304 Hence,  $c \in (f^{m-1} : f_1)$  and  $b = cf \in M_f(f^{m-1} : f_1)$ . The converse is obvious.

305 (3) Let  $b \in (f) \cap (f^m : f_1)$  and  $b' \in (f^m : f_1)$ . Then

306 
$$\left( \frac{f_1 b}{f^m} \right) b' = b \left( \frac{f_1 b'}{f^m} \right) \in (f),$$

307 since  $b \in (f)$ . Hence  $[\langle \frac{f_1 b}{f^m}, b' \rangle]_{\mathbf{B}_0} = 0$  and the bilinear form in (13) vanishes on  $(f) \cap (f^m : f_1)$ .  
 308 Taking the quotient by  $(f) \cap (f^m : f_1)$ , we obtain by part (1) that it is a bilinear form  
 309 defined on  $\tilde{K}_m$  and it has the same expression as Jantzen's form, since  $f^m = t^m$ , so they  
 310 coincide.  $\square$

### 311 3 Transporting the signatures to the algebra A

312 The aim of this section is to establish a relationship between the higher order bilinear forms  
 313  $\langle \cdot, \cdot \rangle_{L_A, f, m}$  (5) and their signatures  $\sigma_{A, f, m}$  in the algebra  $\mathbf{A}$  and Jantzen's relative forms  $\langle \cdot, \cdot \rangle^m$   
 314 (24) and their signatures  $\tau_m$  in  $\mathbf{B}_0$ .

315 Define the isomorphism of  $\mathcal{B}$ -modules

316 
$$\begin{array}{ccc} \mathcal{B} & & \mathcal{B} \\ \cup & & \cup \\ (f : f_1) & \xrightarrow{\Phi} & (f_1 : f) \end{array}$$

317 
$$\Phi(b) = \frac{bf_1}{f}, \quad \Phi^{-1}(c) = \frac{cf}{f_1}$$

318 **Lemma 3.1** *The isomorphism  $\Phi$  induces isomorphisms of  $\mathcal{B}$ -modules, for  $m \geq 1$ :*

319 
$$\Phi : (f^m : f_1) \longrightarrow (f_1 : f) \cap (f^{m-1}) \tag{26}$$
  
 320 
$$\Phi : (f^m) \longrightarrow (f^{m-1} f_1)$$
  
 321 
$$\varphi : \tilde{K}_1 = \text{Ann}_{\mathbf{B}_0}(f_1) \longrightarrow K_1 = \text{Ann}_{\mathbf{A}}(f).$$

322 *Proof* If  $b \in (f^m : f_1)$ , then there exists  $c \in \mathcal{B}$  such that  $bf_1 = cf^m$ . Hence

323 
$$\Phi(b) = \frac{bf_1}{f} = cf^{m-1} \in (f_1 : f) \cap (f^{m-1}).$$

324 Conversely, if  $c = df^{m-1} \in (f_1 : f)$ , then

325 
$$df^m = cf = ef_1 \Rightarrow e = \Phi^{-1}(c) \in (f^m : f_1).$$

326 This proves the first assertion. The second one is just  $\Phi(bf^m) = bf^{m-1} f_1$ . The third assertion  
 327 is obtained by taking the quotient of the first assertion in the Lemma by the second relation  
 328 in the case  $m = 1$ . □

329 Let  $f_2, \dots, f_n$  be a regular sequence of holomorphic functions, denote the volume form  
 330 by  $dVol = dx_1 \wedge \dots \wedge dx_n$ , and let  $\mathbf{Z}_{\mathbb{C}}$  be the complete intersection  $f_2 = \dots = f_n$  as in  
 331 Sect. 2. For any holomorphic function  $g$  define the Jacobian of  $g$  by

332 
$$dg \wedge df_2 \wedge \dots \wedge df_n := Jac(g) dVol, \quad Jac(g) = \begin{vmatrix} \frac{\partial g}{\partial x_1} & \dots & \frac{\partial g}{\partial x_n} \\ \frac{\partial f_2}{\partial x_1} & \dots & \frac{\partial f_2}{\partial x_n} \\ \dots & \dots & \dots \\ \frac{\partial f_n}{\partial x_1} & \dots & \frac{\partial f_n}{\partial x_n} \end{vmatrix}.$$

333 Recall the construction of the generator of the Rosenlicht differentials (see [13]) or dualizing  
 334 module, which is a rational differential form  $\omega_0$  on  $\mathbf{Z}_{\mathbb{C}}$  having the property

335 
$$\omega_0 \wedge (df_2 \wedge \dots \wedge df_n)|_{\mathbf{Z}_{\mathbb{C}}} = dVol|_{\mathbf{Z}_{\mathbb{C}}} \in \frac{\Omega_{\mathbb{C}^n}^n}{(f_2, \dots, f_n)\Omega_{\mathbb{C}^n}^n}.$$

336 The dualizing module on  $\mathbf{Z}_{\mathbb{C}}$  is then  $\mathcal{O}_{\mathbf{Z}_{\mathbb{C}}}\omega_0$ , and it consists of all rational differential forms  
 337  $\sigma$  on  $\mathbf{Z}_{\mathbb{C}}$  that have the property that the residue at 0 of  $h\sigma$  is 0, for any holomorphic function  
 338  $h$  on  $\mathbf{Z}_{\mathbb{C}}$ . Recall also that the residue of a differential form  $\sigma$  at  $0 \in \mathbf{Z}_{\mathbb{C}}$  is the sum of the  
 339 residues of the rational differential form  $\nu^*\sigma$  at  $\nu^{-1}(0)$ , where  $\nu$  is the normalization map  
 340 of  $\mathbf{Z}_{\mathbb{C}}$ . Directly from the definitions above, one obtains that for any holomorphic function  $g$   
 341 on  $\mathbb{C}^n$

342 
$$d(g|_{\mathbf{Z}_{\mathbb{C}}}) = Jac(g)|_{\mathbf{Z}_{\mathbb{C}}}\omega_0.$$

343 Note that the logarithmic derivative of  $g|_{\mathbf{Z}_{\mathbb{C}}}$  is  $\frac{Jac(g)}{g}\omega_0$  and its residue at 0 is the sum of the  
 344 vanishing orders of the function  $g \circ \nu$  at  $\nu^{-1}(0)$ , and hence a positive integer.

345 **Lemma 3.2** *Let  $f, f_1, \dots, f_n$ , and  $\varphi$  be as in Sect. 2. Let  $J_{\mathbf{B}_0}$  and  $J_{\mathbf{A}}$  be the Jacobians of*  
 346  *$(f, f_2, \dots, f_n)$  and  $(f_1, \dots, f_n)$  respectively. Then there exists a positive constant  $c = c(f)$*   
 347 *such that  $\varphi(J_{\mathbf{B}_0}) = cJ_{\mathbf{A}}$ .*

348 *Proof* Since  $([f]_{\mathbf{A}}) \subsetneq \mathbf{A}$ , then taking orthogonal of this relation, we obtain that the ideal  $K_1$   
 349 is not the 0-ideal, and hence  $K_1$  contains the socle. Since  $\varphi : \tilde{K}_1 \rightarrow K_1$  is an isomorphism of  
 350 non-zero ideals, each containing its corresponding 1-dimensional socle, the map  $\varphi$  sends the  
 351 socle ideal to the corresponding socle ideal. Hence  $\varphi$  sends the Jacobian of  $\mathbf{B}_0$  to a non-zero  
 352 multiple of the corresponding Jacobian of  $\mathbf{A}$ .

353 Thus we know that there is a non-zero real number  $c$  with the property

$$354 \quad \left[ \frac{f_1 Jac(f)}{f} \right]_{\mathbf{A}} = c [Jac(f_1)]_{\mathbf{A}}.$$

355 Hence there is a holomorphic function  $h$  on  $\mathbf{Z}_{\mathbb{C}}$  with the property

$$356 \quad \frac{f_1 Jac(f)}{f} - c Jac(f_1) = hf_1$$

357 Dividing by  $f_1$  and multiplying by  $\omega_0$  we obtain

$$358 \quad \frac{Jac(f)}{f} \omega_0 - c \frac{Jac(f_1)}{f_1} \omega_0 = h \omega_0$$

359 Taking residues at 0 we obtain that

$$360 \quad n_1 + \dots + n_r - c(m_1 + \dots + m_r) = 0$$

361 where the  $n_i$  and  $m_j$  are the vanishing orders of the functions  $f \circ \nu$  and  $f_1 \circ \nu$  at  $\nu^{-1}(0)$ ,  
 362 respectively. Hence  $c$  is a positive rational number.  $\square$

363 The real valued bilinear forms on  $\mathbf{A}$  and on  $\mathbf{B}_0$  depended on the choice of real valued  
 364 linear functions  $L_{\mathbf{A}} : \mathbf{A} \rightarrow \mathbb{R}$  and  $L_{\mathbf{B}_0} : \mathbf{B}_0 \rightarrow \mathbb{R}$  which have the property of sending  
 365 the corresponding Jacobians to a positive number. Having chosen  $\mathcal{L}$  and hence  $L_{\mathbf{B}_0}$ , we will  
 366 choose  $L_{\mathbf{A}}$  subject to the compatibility condition

$$367 \quad L_{\mathbf{B}_0}|_{\tilde{K}_1} = L_{\mathbf{A}} \circ \varphi. \tag{27}$$

368 *Proof of Theorem 0.3* Let  $m \geq 1$  and consider the commutative diagram:

$$369 \quad \begin{array}{ccccc} (f^m : f_1) \oplus (f^m : f_1) & \xrightarrow{\frac{f_1 \cdot}{f^m \cdot}} & (f^m : f_1) & \xrightarrow{\tilde{\pi}_0} & \tilde{K}_1 \xrightarrow{L_{\mathbf{B}_0}} \mathbb{R} \\ \Phi \oplus \Phi \downarrow & & \downarrow \Phi & & \downarrow \varphi \quad \downarrow Id. \\ (f_1 : f) \cap (f^{m-1}) \oplus (f_1 : f) \cap (f^{m-1}) & \xrightarrow{\frac{1 \cdot}{f^{m-1} \cdot}} & (f_1 : f) \cap (f^{m-1}) & \xrightarrow{\pi_0} & K_1 \xrightarrow{L_{\mathbf{A}}} \mathbb{R} \end{array}$$

370 Here the mapping  $\frac{f_1 \cdot}{f^m \cdot}$  acts on a couple

$$371 \quad (a, b) \in (f^m : f_1) \oplus (f^m : f_1) \quad \text{by} \quad (a, b) \rightarrow \frac{af_1}{f^m} b,$$

372 and similarly for  $\frac{1 \cdot}{f^{m-1} \cdot}$ . The mapping  $\pi_0$  is obtained by reducing mod  $(f_1)$ . The vertical maps  
 373 are isomorphisms, so we may interpret the commutative diagram as providing a conjugation  
 374 of the top bilinear form into the bottom bilinear form. We reduce the first row by  $(f)$  and the  
 375 second row by  $(f_1)$ . This is possible since  $\Phi(f) = f_1$  and both bilinear forms degenerate  
 376 in the submodules in the denominator of the quotient. We thus obtain that the  $m^{th}$  Jantzen's  
 377 bilinear form is being conjugated by  $\varphi : \tilde{K}_m \rightarrow K_m$  to the bilinear form  $\langle \cdot, \cdot \rangle_{L_{\mathbf{A}}, f, m}$ .  $\square$

378 **4 The index of contact vector fields**

379 4.1 The GSV-index  $Ind_{V_0, \pm}(X|V_0)$  and the signature function  $Sgn_{f,0}(X)$

380 Let  $f : \mathbb{R}^{2N+1} \rightarrow \mathbb{R}$  be a germ of a real analytic function with an algebraically isolated  
 381 singularity at 0. Denote also by  $f$  its extension to a germ in  $\mathbb{C}^{2N+1}$  and let  $V_t$  and  $V_t^{\mathbb{C}}$  be  
 382 the germs of real or complex analytic varieties defined by  $f = t$ . In this section we prove  
 383 Theorem 0.4.

384 We know that both the GSV-index  $Ind_{V_0, \pm}(X|V_0)$  and the signature function  $Sgn_{f,0}(X)$   
 385 verify the law of conservation of numbers (see (29) and similarly for the signature function  
 386  $Sgn_{f,0}(X)$  [6]). They also coincide in smooth points of the variety. Hence, the two indices  
 387 differ by a constant  $K_+$  or  $K_-$  depending only on the function  $f$  (and not on the vector field)  
 388 and on the positive or negative sign chosen in the GSV-index. Given a function  $f$  as above,  
 389 in order to determine these constants  $K_{\pm}$ , it is sufficient to calculate both indices for one  
 390 vector field  $X_0$  tangent to  $V_0$ .

391 *Proof of Theorem 0.4* In order to prove Theorem 0.4, we have to study the index of a family  
 392 of vector fields tangent to the smoothening  $f = t$  of the singular variety  $f = 0$ . When the  
 393 ambient space is even dimensional, this was done [7] using the Hamiltonian vector field  
 394 associated to  $f$ . Here, we study the odd-dimensional ambient space  $\mathbb{R}^{2N+1}$  and we use the  
 395 vector fields

396 
$$X_t = (f - t) \frac{\partial}{\partial x_1} + \sum_{i=1}^N \left[ \frac{\partial f}{\partial x_{2i+1}} \frac{\partial}{\partial x_{2i}} - \frac{\partial f}{\partial x_{2i}} \frac{\partial}{\partial x_{2i+1}} \right],$$

397 which we call the contact vector fields. The vector field  $X_t$  is tangent to  $V_t$ , for any  $t$ , since  
 398  $D(f - t)X_t = \frac{\partial f}{\partial x_1}(f - t)$ , where  $\frac{\partial f}{\partial x_1}$  is the cofactor. For almost all linear hyperplanes  
 399 through 0 in  $\mathbb{C}^{2N+1}$  the projection to this hyperplane gives a description of  $V_0^{\mathbb{C}}$  as a branched  
 400 finite analytic cover [10]. Set  $f_j := \frac{\partial f}{\partial x_j}$ , with  $j = 1, \dots, 2N + 1$ . After perhaps a generic  
 401 rotation, we may assume that 0 is the only point in its neighborhood that satisfies the equations  
 402  $f = f_2 = \dots = f_{2N+1} = 0$ , or equivalently such that  $f, f_2, \dots, f_{2N+1}$  is a regular sequence  
 403 [4]. Hence, the vector field  $X_0$  has an algebraically isolated zero at the origin. The functions  
 404  $f_1, \dots, f_{2N+1}$  form a regular sequence, since  $f$  has isolated singularities. The hypothesis of  
 405 the previous part of this paper are satisfied and we apply Sects. 1, 2 and 3 to this situation.  
 406 Choose a small neighborhood  $U_{\mathbb{C}}$  of  $0 \in \mathbb{C}^{2N+1}$  and a small  $\varepsilon > 0$ , as in Sect. 2.1. The  
 407 derivative of  $X_t$  is

408 
$$DX_t := \begin{pmatrix} f_1 & f_2 & \dots & f_{2N+1} \\ * & \frac{\partial(f_3, -f_2, \dots, f_{2N+1}, -f_{2N})}{\partial(x_2, \dots, x_{2N+1})} \end{pmatrix}. \tag{28}$$

409 Denote by  $Y_t := X_t|_{V_t^{\mathbb{C}}}$  the restriction of  $X_t$  to  $V_t^{\mathbb{C}}$  or to  $V_t$ . The singularities of  $X_t$  are  
 410 always contained in  $V_t^{\mathbb{C}}$ , and hence  $X_t$  and  $Y_t$  have the same singularities:  $\mathbf{Z}_{\mathbb{C}} \cap V_t^{\mathbb{C}}$ .

411 By definition (see [8]), the GSV-index  $Ind_{V_{\pm},0}(Y_0)$  is the sum of the indices of  $Y_t$  at the  
 412 points  $p_t \in V_t, \pm t > 0$  small:

413 
$$Ind_{V_0, \pm}(Y_0, 0) = \sum_{\substack{p_t \in U \cap V_t, Y_t(p_t) = 0 \\ \pm t > 0}} Ind_{V_t}(Y_t, p_t). \tag{29}$$

414 Note that  $V_t$  is smooth, so the signatures  $Ind_{V_t}(Y_t, p_t)$  can be calculated using the usual  
 415 Eisenbud–Levine, Khimshiashvili formula, on the smooth variety  $V_t$ . That is, instead of  
 416 using the Jacobian  $J(X_t)$  as the generator of the socle, one uses the relative Jacobian  $J(Y_t)$ .  
 417 In the localization of the algebra  $\mathbf{B}_t$  in  $p_t$ , we have

$$418 \quad J(X_t) = f_1 J(Y_t). \quad (30)$$

419 Hence, the signature of the bilinear form  $\langle \cdot, \cdot \rangle_t^{rel}$  (10) gives the GSV-index:

$$420 \quad Ind_{V_{\pm,0}}(X_0) = \tau_{\pm}. \quad (31)$$

421 On the other hand, by definition [6], the signature function  $Sgn_{f,0}(X)$  is given by the signature  
 422 of the form  $\langle \cdot, \cdot \rangle_t^{rel}$ , for  $t = 0$ . That is,

$$423 \quad Sgn_{f,0}(X_0) = \tau_0 \quad (32)$$

424 It now follows from Jantzen’s Proposition 2.1 that the constants  $K_{\pm}$  in Theorem 0.4 are

$$425 \quad K_+ = \sum_{m \geq 1} \tau_m, \quad K_- = \sum_{m \geq 1} (-1)^m \tau_m. \quad (33)$$

426 The Theorem 0.4 finally follows from (33) applying Theorem 0.3, which asserts that  $\tau_m =$   
 427  $\sigma_{A,f,m}$ .  $\square$

428 **Corollary 2** Let  $\sigma_A$  be the signature of the Jacobian algebra  $\mathbf{A}$  in (3), and let  $\sigma_{A,f,m}$ ,  $m =$   
 429  $0, \dots, \ell + 1$ , be defined as above. Then

$$430 \quad \sigma_A = \frac{\chi_+ - \chi_-}{2} = \sum_{m=odd} \sigma_{A,f,m}$$

431 *Proof* By Arnold’s formula [1],  $2\sigma_A$  equals  $\chi_+ - \chi_-$ . Now, by the Poincaré–Hopf index  
 432 theorem,  $\chi_+ - \chi_-$  equals  $Ind_{V_{0,+}}(X) - Ind_{V_{0,-}}(X)$ , where  $X$  is a real vector field having  
 433 an algebraically isolated singularity at the origin tangent to  $V$ . The Corollary now follows  
 434 from Theorem 0.4.  $\square$

## 435 4.2 Examples

436 *Example 4.1* Let  $f$  be a quasi-homogeneous real analytic function with an algebraically  
 437 isolated singularity, i.e.  $[f]_{\mathbf{A}} = 0 \in \mathbf{A}$ . In this case,  $Ann_{\mathbf{A}}(f) = \mathbf{A}$ ,  $M_f = 0$  and  $\sigma_{f,1} = \tau_1$   
 438 is the only non-zero Jantzen signature of order higher than 0 and it is equal to  $\sigma_A$ . Hence  
 439  $K_{\pm} = \chi_{\pm} = \pm \sigma_A$ .

440 *Example 4.2* Let  $f = (x^2 + y^3)(x^3 + y^2) + z^2$  and  $V = f^{-1}(0) \subset \mathbb{R}^3$ . This example is  
 441 not a quasi-homogeneous singularity. All calculations have been done using the Computer  
 442 algebra system Singular [9]. The local algebra  $\mathbf{A} = \frac{\mathcal{A}_{\mathbb{R}^3,0}}{(f_x, f_y, f_z)}$  has dimension 11,  $Ann_{\mathbf{A}}(f)$  is  
 443 the maximal ideal of dimension 10 and  $([f]_{\mathbf{A}})$  is the 1-dimensional socle ideal. We thus have  
 444 that  $M_f$  has nine one-dimensional Jordan blocks and one two-dimensional Jordan block. The  
 445 Hessian  $[Hess(f)]_{\mathbf{A}}$  generating the socle equals  $-220[f]_{\mathbf{A}}$  in  $\mathbf{A}$ . The filtration (4) is given  
 446 by

$$447 \quad (f) \subset Ann_{\mathbf{A}}(f) \subset \mathbf{A}.$$

448 The signature  $\sigma_1$  is the signature of  $\langle \cdot, \cdot \rangle_{L_{\mathbf{A}}}$  on the 9 dimensional space isomorphic to  $\frac{Ann_{\mathbf{A}}(f)}{(f)}$ .

449 This signature is equal to 1. The signature  $\sigma_2$  is given by the the sign of  $L_{\mathbf{A}}(\frac{f \cdot f}{f}) = L_{\mathbf{A}}(f) <$   
 450  $0$ , so  $\sigma_2 = -1$ . This gives by Theorem 2 that  $K_+ = 0$ ,  $K_- = -2$ .

451 **References**

- 452 1. Arnold, V.: Index of a singular point of a vector field, the Petrovskii–Oleinik inequality and mixed Hodge  
453 structures. *Funct. Anal. Appl.* **12**(1), 1–14 (1978)
- 454 2. Bonatti, Ch., Gómez-Mont, X.: The index of a holomorphic vector field on a singular variety, I. *Asté-*  
455 *risque* **222**, 9–35 (1994)
- 456 3. Eisenbud, D., Levine, H.: An algebraic formula for the degree of a  $C^\infty$  map germ. *Ann. Math.* **106**, 19–38  
457 (1977)
- 458 4. Eisenbud, D.: *Commutative Algebra with a View Towards Algebraic Geometry*. Springer, Heidelberg  
459 (1995)
- 460 5. Giraldo, L., Gómez-Mont, X., Mardešić, P.: Computation of topological numbers via linear algebra:  
461 Hypersurfaces, vector fields and vector fields on hypersurfaces. *Contemp. Math.* **240**, 175–182 (1999)
- 462 6. Gómez-Mont, X., Mardešić, P.: The index of a vector field tangent to a hypersurface and the signature of  
463 the relative Jacobian determinant. *Ann. Inst. Fourier* **47**(5), 1523–1539 (1997)
- 464 7. Gómez-Mont, X., Mardešić, P.: The index of a vector field tangent to an odd dimensional hypersurface  
465 and the signature of the relative Hessian. *Funct. Anal. Appl.* **33**(1) (1999)
- 466 8. Gómez-Mont, X., Seade, J., Verjovsky, A.: The index of a holomorphic flow with an isolated singula-  
467 rity. *Math. Ann.* **291**, 737–751 (1991)
- 468 9. Greuel, G.-M., Pfister, G., Schönemann, H.: *Singular 3.0. A Computer Algebra System for Polynomial*  
469 *Computations*. Center for Computer Algebra, University of Kaiserslautern (2005). [http://www.singular.](http://www.singular.uni-kl.de)  
470 [uni-kl.de](http://www.singular.uni-kl.de)
- 471 10. Gunning, R.C.: *Introduction to Holomorphic Functions of Several Complex Variables*, vol. II. Local  
472 theory. Wadsworth & Brooks/Cole (1990)
- 473 11. Jantzen, J.C.: *Moduln mit einem Höchsten Gewicht*. *Lecture Notes in Mathematics*, vol. 750. Springer,  
474 Berlin (1979)
- 475 12. Khimshiashvili, G.N.: On the local degree of a smooth map. *Trudi Tbilisi math. Inst.*, pp. 105–124 (1980)
- 476 13. Montaldi, J., van Straten, D.: One-forms on singular curves and the topology of real curves singulari-  
477 ties. *Topology* **29**(4), 501–510 (1990)
- 478 14. Vogan, D.: Unitarizability of certain series of representations. *Ann. Math. (2)* **120**(1), 141–187 (1984)